



(12) **United States Patent**
Wright et al.

(10) **Patent No.:** **US 9,861,582 B2**
(45) **Date of Patent:** ***Jan. 9, 2018**

(54) **PHARMACEUTICAL FORMULATION**
CONTAINING GELLING AGENT

(71) Applicant: **Purdue Pharma L.P.**, Stamford, CT (US)

(72) Inventors: **Curtis Wright**, Rockport, MA (US); **Benjamin Oshlack**, Boca Raton, FL (US); **Christopher Breder**, Bethesda, MD (US)

(73) Assignees: **Purdue Pharma L.P.**, Stamford, CT (US); **The P.F. Laboratories, Inc.**, Totowa, NJ (US); **Purdue Pharmaceuticals L.P.**, Wilson, NC (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **15/015,769**

(22) Filed: **Feb. 4, 2016**

(65) **Prior Publication Data**

US 2016/0151358 A1 Jun. 2, 2016

Related U.S. Application Data

(63) Continuation of application No. 14/638,685, filed on Mar. 4, 2015, which is a continuation of application (Continued)

(51) **Int. Cl.**

A61K 47/36 (2006.01)
A61K 45/06 (2006.01)
A61K 31/485 (2006.01)
A61K 9/20 (2006.01)
A61K 9/00 (2006.01)
A61K 31/439 (2006.01)
A61K 47/38 (2006.01)
A61K 47/10 (2017.01)
A61K 31/167 (2006.01)

(Continued)

(52) **U.S. Cl.**

CPC **A61K 9/2054** (2013.01); **A61J 3/10** (2013.01); **A61K 9/0002** (2013.01); **A61K 9/006** (2013.01); **A61K 9/0053** (2013.01); **A61K 9/06** (2013.01); **A61K 9/08** (2013.01); **A61K 9/1635** (2013.01); **A61K 9/1641** (2013.01); **A61K 9/1652** (2013.01); **A61K 9/20** (2013.01); **A61K 9/205** (2013.01); **A61K 9/2009** (2013.01); **A61K 9/2013** (2013.01); **A61K 9/2018** (2013.01); **A61K 9/2027** (2013.01); **A61K 9/2031** (2013.01); **A61K 9/2095** (2013.01); **A61K 9/28** (2013.01); **A61K 9/284** (2013.01); **A61K 9/2806** (2013.01); **A61K 9/2813** (2013.01); **A61K 9/2846** (2013.01); **A61K 9/2853** (2013.01); **A61K 9/2866** (2013.01); **A61K 9/2893** (2013.01);

A61K 9/48 (2013.01); **A61K 9/485** (2013.01); **A61K 9/4808** (2013.01); **A61K 9/4833** (2013.01); **A61K 9/4858** (2013.01); **A61K 9/4866** (2013.01); **A61K 9/4891** (2013.01); **A61K 9/5047** (2013.01); **A61K 9/5078** (2013.01); **A61K 9/5089** (2013.01); **A61K 9/70** (2013.01); **A61K 31/137** (2013.01); **A61K 31/167** (2013.01); **A61K 31/192** (2013.01); **A61K 31/439** (2013.01); **A61K 31/4458** (2013.01); **A61K 31/48** (2013.01); **A61K 31/485** (2013.01); **A61K 45/06** (2013.01); **A61K 47/02** (2013.01); **A61K 47/08** (2013.01); **A61K 47/10** (2013.01); **A61K 47/12** (2013.01); **A61K 47/14** (2013.01); **A61K 47/26** (2013.01); **A61K 47/32** (2013.01); **A61K 47/34** (2013.01); **A61K 47/36** (2013.01); **A61K 47/38** (2013.01)

(58) **Field of Classification Search**

None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,065,143 A 11/1962 Christenson et al.
3,133,132 A 5/1964 Loeb et al.

(Continued)

FOREIGN PATENT DOCUMENTS

EP 0318262 5/1989
EP 0661045 5/1995

(Continued)

OTHER PUBLICATIONS

U.S. Appl. No. 13/890,874, filed May 9, 2013.
U.S. Appl. No. 14/610,156, filed Jan. 30, 2015.
U.S. Appl. No. 14/658,285, filed Mar. 16, 2015.
U.S. Appl. No. 14/851,575, filed Sep. 11, 2015.
Ansel, Howard C., et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 7th Edition, 1999, pp. 1-2, 23-163, 179-243, 397-449, 552-562, Lippincott Williams & Wilkins, United States.
Apicella, A., "Poly(ethylene oxide)(PEO) and Different Molecular Weight PEO Blends Monolithic Devices for Drug Release," *Biomaterials*, vol. 14, No. 2, 1993, pp. 83-90.
Apicella, et al., "Poly(ethylene oxide)-Based Delivery Systems," *Polymeric Drugs and Drug Administration*, ACS Symposium Series 545, Chapter 9 (1994), pp. 111-125.

(Continued)

Primary Examiner — Robert A Wax
Assistant Examiner — Randeep Singh

(74) *Attorney, Agent, or Firm* — Lowenstein Sandler LLP

(57) **ABSTRACT**

Disclosed in certain embodiments is a controlled release oral dosage form comprising a therapeutically effective amount of a drug susceptible to abuse together with one or more pharmaceutically acceptable excipients; the dosage form further including a gelling agent in an effective amount to impart a viscosity unsuitable for administration selected from the group consisting of parenteral and nasal administration to a solubilized mixture formed when the dosage form is crushed and mixed with from about 0.5 to about 10 ml of an aqueous liquid; the dosage form providing a therapeutic effect for at least about 12 hours when orally administered to a human patient.

22 Claims, No Drawings